

J Infect Dis. 2001 Apr 1;183(7):1071-8. Epub 2001 Mar 8.

Related Articles, Links



**Further characterization of *Escherichia coli* brain microvascular endothelial cell invasion gene *ibeA* by deletion, complementation, and protein expression.**

**Huang SH, Wan ZS, Chen YH, Jong AY, Kim KS.**

Childrens Hospital Los Angeles and University of Southern California Keck School of Medicine, Los Angeles, CA 90027, USA. shhuang@hsc.usc.edu

The *ibeA* gene (*ibe10*) previously identified by *TnphoA* mutagenesis is part of a 50-kDa full-length open-reading frame (ORF) encoded by a 1.37-kb DNA fragment. An isogenic in-frame deletion mutant of *ibeA* (ZD1) was constructed by chromosomal gene replacement with a suicide plasmid pCVD442 carrying a 2.1-kb DNA fragment with an *ibeA* deletion. Similar to the previously described *TnphoA* insertion mutant of *ibeA*, the isogenic *ibeA* deletion mutant ZD1 was significantly less invasive in human brain microvascular endothelial cells (BMECs) than the parent strain. The mutant ZD1 was fully complemented by the *ibeA* ORF. The *ibeA* gene was subcloned into pET28a(+) and was expressed as a recombinant protein with an N-terminal histidine tag. The recombinant IbeA protein had much greater activity (50 times) in blocking the invasion of BMECs by *Escherichia coli* K1 than did the partial protein fragment, which provides further evidence that *ibeA* is an important determinant for *E. coli* K1 invasion of BMECs.

PMID: 11237832 [PubMed - indexed for MEDLINE]



A service of the National Library of Medicine  
and the National Institutes of Health

My NCBI

[\[Sign In\]](#) [\[Regis\]](#)

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search



for



Limits

Preview/Index

History

Clipboard

Details

Display



Show



Sort by



Send to



About Entrez  
NCBI Toolbar

All: 649 Review: 35

Text Version

Items 1 - 20 of 649

Page

of 33 Next

Entrez PubMed  
Overview  
Help | FAQ  
Tutorials  
New/Noteworthy  
E-Utilities

PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
Special Queries  
LinkOut  
My NCBI

Related Resources  
Order Documents  
NLM Mobile  
NLM Catalog  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

- ☐ 1: [Huang SH, Chen YH, Kong G, Chen SH, Besemer J, Borodovsky M, Jong A.](#) Related Articles, Links



A novel genetic island of meningitic Escherichia coli K1 containing the ibeA invasion gene (GimA): functional annotation and carbon-source-regulated invasion of human brain microvascular endothelial cells.

Funct Integr Genomics. 2001 Sep;1(5):312-22.

PMID: 11793250 [PubMed - indexed for MEDLINE]

- ☐ 2: [Huang SH, Chen YH, Fu Q, Stins M, Wang Y, Wass C, Kim KS.](#) Related Articles, Links



Identification and characterization of an Escherichia coli invasion gene locus, ibeB, required for penetration of brain microvascular endothelial cells.

Infect Immun. 1999 May;67(5):2103-9.

PMID: 10225861 [PubMed - indexed for MEDLINE]

- ☐ 3: [Huang SH, Wass C, Fu Q, Prasadarao NV, Stins M, Kim KS.](#) Related Articles, Links



Escherichia coli invasion of brain microvascular endothelial cells in vitro and in vivo: molecular cloning and characterization of invasion gene ibe10.

Infect Immun. 1995 Nov;63(11):4470-5.

PMID: 7591087 [PubMed - indexed for MEDLINE]

- ☐ 4: [Huang SH, Wan ZS, Chen YH, Jong AY, Kim KS.](#) Related Articles, Links



Further characterization of Escherichia coli brain microvascular endothelial cell invasion gene ibeA by deletion, complementation, and protein expression.

J Infect Dis. 2001 Apr 1;183(7):1071-8. Epub 2001 Mar 8.

PMID: 11237832 [PubMed - indexed for MEDLINE]

- ☐ 5: [Wang Y, Huang SH, Wass CA, Stins MF, Kim KS.](#) Related Articles, Links



The gene locus yijP contributes to Escherichia coli K1 invasion of brain microvascular endothelial cells.

Infect Immun. 1999 Sep;67(9):4751-6.

PMID: 10456927 [PubMed - indexed for MEDLINE]

- ☐ 6: [Badger JL, Wass CA, Kim KS.](#) Related Articles, Links



Identification of Escherichia coli K1 genes contributing to human brain microvascular endothelial cell invasion by differential fluorescence induction.

Mol Microbiol. 2000 Apr;36(1):174-82.

PMID: 10760174 [PubMed - indexed for MEDLINE]

☐ 7: [Wang Y, Kim KS.](#)[Related Articles, Links](#)

Effect of rpoS mutations on stress-resistance and invasion of brain microvascular endothelial cells in Escherichia coli K1.

FEMS Microbiol Lett. 2000 Jan 15;182(2):241-7.

PMID: 10620673 [PubMed - indexed for MEDLINE]

☐ 8: [Wang Y, Kim KS.](#)[Related Articles, Links](#)

Role of OmpA and IbeB in Escherichia coli K1 invasion of brain microvascular endothelial cells in vitro and in vivo.

Pediatr Res. 2002 May;51(5):559-63.

PMID: 11978877 [PubMed - indexed for MEDLINE]

☐ 9: [Hoffman JA, Badger JL, Zhang Y, Huang SH, Kim KS.](#)[Related Articles, Links](#)

Escherichia coli K1 aslA contributes to invasion of brain microvascular endothelial cells in vitro and in vivo.

Infect Immun. 2000 Sep;68(9):5062-7.

PMID: 10948126 [PubMed - indexed for MEDLINE]

☐ 10: [Prasadarao NV, Wass CA, Huang SH, Kim KS.](#)[Related Articles, Links](#)

Identification and characterization of a novel Ibe10 binding protein that contributes to Escherichia coli invasion of brain microvascular endothelial cells.

Infect Immun. 1999 Mar;67(3):1131-8.

PMID: 10024553 [PubMed - indexed for MEDLINE]

☐ 11: [Badger JL, Wass CA, Weissman SJ, Kim KS.](#)[Related Articles, Links](#)

Application of signature-tagged mutagenesis for identification of escherichia coli K1 genes that contribute to invasion of human brain microvascular endothelial cells.

Infect Immun. 2000 Sep;68(9):5056-61.

PMID: 10948125 [PubMed - indexed for MEDLINE]

☐ 12: [Khan NA, Shin S, Chung JW, Kim KJ, Elliott S, Wang Y, Kim KS.](#)[Related Articles, Links](#)

Outer membrane protein A and cytotoxic necrotizing factor-1 use diverse signaling mechanisms for Escherichia coli K1 invasion of human brain microvascular endothelial cells.

Microb Pathog. 2003 Jul;35(1):35-42.

PMID: 12860457 [PubMed - indexed for MEDLINE]

☐ 13: [Prasadarao NV, Wass CA, Kim KS.](#)[Related Articles, Links](#)

Endothelial cell GlcNAc beta 1-4GlcNAc epitopes for outer membrane protein A enhance traversal of Escherichia coli across the blood-brain barrier.

Infect Immun. 1996 Jan;64(1):154-60.

PMID: 8557333 [PubMed - indexed for MEDLINE]

☐ 14: [Kim KS.](#)[Related Articles, Links](#)

E. coli invasion of brain microvascular endothelial cells as a pathogenetic basis of meningitis.

Subcell Biochem. 2000;33:47-59. Review.

PMID: 10804851 [PubMed - indexed for MEDLINE]

☐ 15: [Teng CH, Cai M, Shin S, Xie Y, Kim KJ, Khan NA, Di Cello F.](#)[Related Articles, Links](#)

Kim KS.



Escherichia coli K1 RS218 interacts with human brain microvascular endothelial cells via type 1 fimbria bacteria in the fimbriated state.

Infect Immun. 2005 May;73(5):2923-31.

PMID: 15845498 [PubMed - indexed for MEDLINE]

- ☐ 16: Prasadarao NV, Wass CA, Weiser JN, Stins MF, Huang SH, Kim KS. [Related Articles](#), [Links](#)



Outer membrane protein A of Escherichia coli contributes to invasion of brain microvascular endothelial cells.

Infect Immun. 1996 Jan;64(1):146-53.

PMID: 8557332 [PubMed - indexed for MEDLINE]

- ☐ 17: Khan NA, Wang Y, Kim KJ, Chung JW, Wass CA, Kim KS. [Related Articles](#), [Links](#)



Cytotoxic necrotizing factor-1 contributes to Escherichia coli K1 invasion of the central nervous system.

J Biol Chem. 2002 May 3;277(18):15607-12. Epub 2002 Feb 27.

PMID: 11877402 [PubMed - indexed for MEDLINE]

- ☐ 18: Hoffman JA, Badger JL, Zhang Y, Kim KS. [Related Articles](#), [Links](#)



Escherichia coli K1 purA and sorC are preferentially expressed upon association with human brain microvascular endothelial cells.

Microb Pathog. 2001 Aug;31(2):69-79.

PMID: 11453702 [PubMed - indexed for MEDLINE]

- ☐ 19: Wang Y, Wen ZG, Kim KS. [Related Articles](#), [Links](#)



Role of S fimbriae in Escherichia coli K1 binding to brain microvascular endothelial cells in vitro and penetration into the central nervous system in vivo.

Microb Pathog. 2004 Dec;37(6):287-93. Epub 2004 Dec 9.

PMID: 15619424 [PubMed - indexed for MEDLINE]

- ☐ 20: Bloch CA, Huang SH, Rode CK, Kim KS. [Related Articles](#), [Links](#)



Mapping of noninvasion TnphoA mutations on the Escherichia coli O18:K1:H7 chromosome.

FEMS Microbiol Lett. 1996 Nov 1;144(2-3):171-6.

PMID: 8900061 [PubMed - indexed for MEDLINE]

Items 1 - 20 of 649

Page

1

of 33 Next

Display

Summary



Show

20



Sort by



Send to



[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Dec 22 2005 16:39:56

0686] Gormelen A, Blondelet-Rouault M H, Pernodet J L. Characterization of a glycosyl transferase inactivating macrolides, encoded by gimA from *Streptomyces ambofaciens*. Antimicrob Agents Chemother. (1998). 42(10): 2612-9.

Analysis of the protein sequence deduced from orf29, using the SignalP program (<http://www.cbs.dtu.dk/services/SignalP/>) (Nielsen, H., et al., 1997), shows that this protein has a C-terminal signal sequence with a predicted cleavage site between positions 30 and 31 (QSA/QA). It may be predicted that this protein is extracellular. It might, as a glycosylhydrolase, have a role in the reactivation of spiramycin inactivated by glycosylation by the glycosyltransferases GimA and/or GimB (Gourmelen et al, 1998).